

---

# Water Column Evaluation

**Alan J Kennedy**

**Research Biologist**

**US Army ERDC, Vicksburg, MS**

**Email: [Alan.J.Kennedy@usace.army.mil](mailto:Alan.J.Kennedy@usace.army.mil)**





---

**Dredged Material Assessment and Management Seminar  
15-17 September 2009, Detroit, MI**



# Water Column Evaluation (Conceptual Model)

Potential of DM disposal to cause adverse effects on water column organisms



**Water column toxicity is one assessment considered in DM disposal**


# Water Column Evaluation

## Approach: Open water disposal of Dredged Material


---

- **Main discussion points**
  - **DM is suspended in water for a short period**
  - **Short-term water column exposure and effects**
  - **Can a factual determination be made from existing information (chemical, toxicity values)?**
    - **Relate to applicable water quality standards**
    - **If more information needed, conduct bioassays**


# Water Column Evaluation




Tiered process → follow as far as necessary to make decision



# Water Column Evaluation (Decision Tree)



# Water Column Evaluation (Physical / Chemical Testing)



## Contaminant concentration in disposed DM:

- **Ocean disposal (Ocean Testing Manual)**
  - Seaward of national baseline
  - Marine Protection, Research and Sanctuaries Act (MPRSA)
  - Limiting Permissible Concentration (LPC)
    - **Definition:** Water concentration not to be exceeded by DM constituents after mixing
    - Based on Water Quality standards, or
    - An acute LC50 adjusted by an application factor (usually 0.01)
- **Inland disposal (Inland Testing manual)**
  - Landward of national baseline, rivers, lakes
  - Clean Water Act
  - Mixing zones variable – contingent on state, water body
  - Compliance with WQS (at least as strict as national standards)






Figure 1-1. Geographical Jurisdictions of the MPRSA and CWA  
From USEPA / USACE. 2004. EPA842-B-92-008.

# Water Column Evaluation (Physical / Chemical Testing)




Must meet LPC after  
4 hours mixing

Must meet LPC/WQS at all times



# TIER II: Two step process



## 1. Screening Step:

- Conduct chemical analysis of DM for CoCs
- Make very conservative assumption
  - 100% DM contaminants goes to water
- For contaminant requiring greatest dilution (D):
  - DM < LPC or WQS → DM complies → **STOP**
  - DM > LPC or WQS → **Move to step 2**

$$D = C_s * \frac{SS}{1000} - \frac{C_{wq}}{C_{wq} - C_{ds}}$$

D = Dilution to meet WQS and / or WQC  
 $C_s$  = contaminant concentration in the sediment  
 SS = suspended solids concentration  
 $C_{wq}$  = WQS and / or WQC  
 $C_{ds}$  = Disposal site concentration

## 2. Elutriate preparation step:

- More realistic chemical analysis
- Use more representative dissolved concentrations in mixing model
- No biological testing



$$D = \frac{C_e - C_{wq}}{C_{wq} - C_{ds}}$$

$C_e$  = concentration of the dissolved contaminant in the standard elutriate

Compare above values to LPC / WQS  
 Apply data into predictive numerical mixing model (Appendix C)



# TIER II: Step Two: Prepare Elutriate





1. Remove overlying water
2. Centrifugation / filtration
3. Chemical analysis



Media Type	Application
Dredged Material (1 part)	Elutriate preparation
Dredging Site Water (4 parts)	Elutriate preparation

# TIER II: Possible conclusions



## 1. DM exceeds LPC / WQS

- ✓ Needs management action
- ✓ No further testing needed


## 2. DM meets LPC / WQS:

- ✓ 4-hours within mixing zone (MPRSA)
- ✓ At all times outside mixing zone

## 3. DM meets LPC / WQS but...


- ✓ WQC not available some contaminant(s)
- ✓ Concern for contaminant interactions
  - Move to Tier III analysis

# TIER III: Overview




- **Biological testing conducted if Tiers I / II:**
  - Inadequate information for factual determination
  - CoCs that lack WQS
  - Contaminants present at potentially adverse levels (gray area)
  - Potential for unknown chemicals or interactions
- **Tier III**
  - Biological exposures conducted
  - Evaluate potential for toxicity
    - Generate lethal/effective median concentration (L(E)C50)
    - Relate toxicity information to mixing model / standards

# TIER III: Biological Testing Summary




- Prepare elutriate (as before)



- Remove overlying water
- Contains both dissolved and suspended DM
  - Centrifuge / filter
    - Only if necessary
- Assess survival across elutriate dilution
- Apply resulting toxicity data to mixing model


Media Type	Application
Dredged Material (1 part)	Elutriate preparation
Dredging Site Water (4 parts)	Elutriate preparation
Disposal Site Water (preferred)	Dilution of elutriate Reference water
Reconstituted Water (other approved water)	

# TIER III: Test Species Selection



- **Three species of different phyla recommended to evaluate the potential for elutriate toxicity**
  - Zooplankton, crustaceans, fish, molluscs, (phytoplankton)
  - MPRSA → must test three species
  - CWA → should test multiple species
  - At least one needs to be a recommended species (previously “benchmark”)
    - Routinely utilized
    - Proven track record
    - National guidance or RIM


# TIER III: Test Species Selection



- **Other test species**
  - Represent organisms indigenous to the disposal site
  - Locally important
  - Regional Implementation Manuals
- **Species selection considerations**
  - Ecological relevance / indigenous
  - Appropriate chemical sensitivity / age class (e.g., larvae, juveniles)
  - Availability of standardized protocol / consistent track record
  - Susceptibility to confounding factors (DO<sub>2</sub>, laboratory handling)
  - Availability year round

# Tier III: Test Species

## Freshwater disposal




- **Freshwater (< 1 ‰)**

- **Arthropoda / Crustacea**

- **Cladocerans (i.e., zooplankton)**


- *Daphnia magna / pulex* \*
- *Ceriodaphnia dubia* \*



- **Vertebrata**

- **Fish**

- *Pimephales promelas* \*
- *Lepomis macrochirus*
- *Oncorhynchus mykiss* \*




OK Dept Wildlife Conservation

\* Recommended species

# Tier III: Test Species

## Marine/estuarine disposal



- **Marine (> 25 ‰)**

- **Echinodermata**

- Urchins, *Strongylocentrotus*, *Arbacia*
    - Sand Dollar, *Dendraster* spp.



- **Arthropoda / crustacea**


- **Shrimp**
      - *Americamysis bahia* \*
      - *Neomysis* \*
      - *Holmesimysis* spp. \*
    - Copepods, *Acartia* sp. \*



- **Estuarine / Marine (1 – 25+ ‰)**

- **Bivalve Molluscs**

- Oysters, *Crassostrea* spp. \*
    - Mussels, *Mytilus* spp. \*



- **Vertebrata**


- Silversides, *Menidia* \*
    - *Cyprinodon variegatus* \*




\* Recommended species



# TIER III: Conduct of Bioassays




<b>Test methods</b>	<b>ITM Appendix E</b>
<b>Exposure</b>	<b>48 or 96-hours</b>
<b>Primary endpoint</b>	<b>Survival or development</b>
<b>Dilutions</b>	<b>Three (10, 50, 100%)</b>
<b>Replicates / dilution</b>	<b>Five</b>
<b>Organisms / replicate</b>	<b>Usually 10</b>
<b>Acceptability criterion</b>	<ul style="list-style-type: none"> <li>• <b>≥70 or 90 % survival</b></li> <li>• <b>Reference toxicity test within range</b></li> </ul>





## Specific testing protocols


- ITM Appendix E
- US EPA / ASTM citations within



# TIER III: Data analysis



# TIER III: Data Analysis (Step 1)





- Survival in undiluted elutriate reduced more than 10% relative to the control?
- Statistical reduction of survival in the undiluted elutriate relative to the control (dilution water)?

Undiluted Elutriate  = 20 ± 8% Survival 


Dilution water (control)  = 90 ± 5% Survival 

- Next step: determine LC50 value, LPC and modeled dilution


# TIER III: Data Interpretation



# TIER III: Data Analysis (Step 2)



Determine the LC50 value




$$\text{LC50 (42\%)} \times 0.01 \text{ (LPC)} = 0.42\%$$




Model output indicates DM is < 0.1% inside and outside the mixing zone

- DM diluted to lower concentration (0.1%) than LPC (0.42%)
- DM elutriate does not exceed LPC / WQS (“passes”)


# Application Factors Considerations



- **NAS (1972): Default = 0.01**
- **EPA: 40cfr 227.29 (3)**
  - Use different AF with scientific rationale
  - $ACR = LC50 / NOEC$ ,  $AF = 1 / ACR$
  - $AF = 0.1$  to  $0.01$  (Verma 1981)
  - 90<sup>th</sup> ACR: 73 ( $AF = 0.01$ ) (Lange 1998)
  - Fish  $AF = 0.15$  (Arsenic) (Lima 1984)
  - $AF = 0.1 \rightarrow 60\%$  of fish (Heger 1995)
  - $AF = 0.01 \rightarrow 90\%$  of fish
- **AF is chemical class specific**
  - Persistent:  $AF = 0.01$
  - Non persistent (half life <8 wks):
    - $AF = 0.05$  to  $0.1$
    - Ammonia:  $AF = 0.11$  (Thurston 1986)




# TIER III: Possible conclusions



- 1. DM discharge toxicity not predicted relative to the reference condition**
- 2. DM discharge toxicity is predicted relative to the reference condition**
- 3. Further information needed for actual determinations**
  - **Move to Tier IV (less common)**

# TIER IV: Case-specific (laboratory / field testing)




- Lower tiers do not provide enough information for factual determinations
  - Rare occasions
  - Inconclusive test results
  - Conflicting evidence
  - Ammonia toxicity suspected
- Specific studies may include:
  - Use of different test species / exposure durations / endpoints (e.g., growth, reproduction)
  - Laboratory *or in situ* exposures (field)
  - TRE/TIE to discriminate ammonia, metals and organic toxicity




# Confounder: Ammonia

## Methods for removal in bioassays




- Algae: *Olva* (Marine)
- pH modifications
  - Increase pH (10) and aerate → decrease pH → test
  - PROBLEM: alteration of metals speciation
- Zeolite Column Treatment (freshwater)
  - Zeolite removes ammonia and metals toxicity
  - EDTA treatment to remove metals
  - PROBLEM: Cannot completely rule out metals toxicity



# Synthesis: Water Column

- **Goal:** Evaluate potential of DM to cause adverse effects on water column organisms
- This is just one pathway to establish a weight of evidence
- Still need to consider other pathways (e.g., benthic effects)
- Generate data to estimate toxicity potential of DM disposal



- **Procedure:** Follow tiered process only as far as necessary to make risk-based determination

#### References cited

- Fava JA, McCulloch WL et al. 1984. Marine Technology Society, Washington DC, 10-12 Sept 1984
- Giesy JP, Graney RL. 1989. *Hydrobiologia* 188/189: 21-60.
- Heger W, Jung SJ, Martin S, Peter H. 1995. *Hydrobiologia* 31: 2707 – 2726.
- Lange R, Hutchinson TH, Scholz N, SolbeJ. 1998. *Chemosphere* 36: 115–127.
- Lima AR, Curtis C, Hammermeister DE et al. 1984. *Arch Environ Contam Toxicol* 13:59 –601.
- Thurston RV, Russo RC, Meyn EL, Zajdel RK, Smith CE. 1986. *Trans Am Fish Soc* 115:196-207.
- Verma SR, Tonk IP, Dalela RC. 1981. *Acta Hydrochim Hydrobiol* 9(3): 247-254.