

Room for the River and Cyclic Floodplain Rejuvenation



Martin Baptist

19 January 2011



Problem background

- Two main problems in the River Rhine are (i) flood safety and (ii) ecological degradation.
- In the Netherlands, more room for the rivers is created in order to increase their conveyance capacity.
- At the same time restoration of natural values in river floodplains is realised.



19 January 2011



5



Conflicting goals

- The combination of nature rehabilitation and flood safety is conflicting.
- Natural growth of floodplain vegetation increases the hydraulic roughness and the sedimentation rate of the floodplains¹.
- Therefore there is an increasing interest in the interactions between vegetation, hydraulics, sediment transport and geomorphology.

¹ Baptist, M.J., et al. (2004). Assessment of Cyclic Floodplain Rejuvenation on Flood Levels and Biodiversity in the Rhine River. *River Research and Applications* 20(3), 285-297.

Open channel flow

- Chézy equation: $u = C\sqrt{hi}$
 - u = depth-averaged flow velocity (m/s)
 - h = water depth (m)
 - i = slope of the bed (-)
 - C = Chézy coefficient ($\text{m}^{1/2} \cdot \text{s}^{-1}$)
- Colebrook-White equation: $C = 18 \log \frac{12R}{k}$ $C = \frac{R^{1/6}}{n}$
 - k = Nikuradse equivalent sand roughness height (m)



Nikuradse roughness height of grassed floodplains

Nikuradse roughness height of grassed floodplains

- Klaassen & Van der Zwaard (1974): 0.07 m
- Klaassen & Van Urk (1985): 0.2 m in winter, 0.7 m in summer.
- RWS (1997): 0.2 m for smooth grassland, 0.5 m for normal grassland, 1.0 m for 'rough' grassland.
- RWS (2003): 0.26 m (at 4 m water depth).
- Baptist (2003): 1.0 m (at 4 m water depth).
- VNK (2006): 0.3 m.
- Straatsma (UU, 2006): 0.07 m.

About 2/3rd of the Rhine floodplains consists of grassland

Natural vegetation succession

- In rivers confined by dikes, vegetation development can only go one way: succession to older stages. Natural processes to remove older vegetation, such as erosion or ice scour, can not occur anymore.



19 January 2011


11

The Allier River (France)

Cyclic Floodplain rejuvenation

- Cyclic Floodplain Rejuvenation (CFR) is the recurrent removal of vegetation and/or sediment,
 - (i) to guard flood safety and
 - (ii) to increase biodiversity.

CFR in Dutch floodplains



Cyclic floodplain destruction?



Conclusions

- Vegetation succession in rivers can diminish the effects of flood protection measures.
- Introducing large herbivores slows down vegetation succession.
- However, floodplain vegetation may still form the bottleneck for flood protection.
- Cyclic Floodplain Rejuvenation is a strategy that might solve the dilemma between flood protection and nature rehabilitation.